Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 5 February 2008 (MN LaTeX style file v2.2)

arXiv:astro-ph/0604235v1 11 Apr 2006

Stochastic wake field particle acceleration in Gamma-Ray Bursts

G.Barbiellini[1], F.Longo[1], N.Omodei[2], A.Celotti[3] and M.Tavani[4,5]
[1]*Dipartimento di Fisica, Università di Trieste and INFN, Sezione di Trieste, via Valerio 2, 34100, Trieste, Italy*
[2]*INFN, Sezione di Pisa, Edificio C - Polo Fibonacci - Largo B. Pontecorvo 3, Pisa, Italy*
[3]*SISSA, via Beirut 2, 34100, Trieste, Italy*
[4]*INAF-IASF, via Fosso del Cavaliere 100, Roma, Italy*
[5]*Dipartimento di Fisica, via della Ricerca Scientifica, Università Tor Vergata, Roma (Italy)*

Received / Accepted

ABSTRACT
Gamma-Ray Burst (GRB) prompt emission can, for specific conditions, be so powerful and short-pulsed to strongly influence any surrounding plasma. In this paper, we briefly discuss the possibility that a very intense initial burst of radiation produced by GRBs satisfy the intensity and temporal conditions to cause stochastic *wake-field particle acceleration* in a surrounding plasma of moderate density. Recent laboratory experiments clearly indicate that powerful laser beam pulses of tens of femtosecond duration hitting on target plasmas cause efficient particle acceleration and betatron radiation up to tens of MeV . We consider a simple but realistic GRB model for which particle wake-field acceleration can first be excited by a very strong low-energy precursor, and then be effective in producing the observed prompt X-ray and gamma-ray GRB emission. We also briefly discuss some of the consequences of this novel GRB emission mechanism.

Key words: gamma-rays: bursts, plasmas, acceleration of particles, radiation mechanisms: non-thermal



Figure 1. Oscillation and radiation produced by a relativistic electron oscillating in an ion channel

1 INTRODUCTION

Gamma-Ray Bursts (GRBs) are major explosive phenomena of our Universe in need of an explanation. The prompt GRB emission in the hypothesis of beamed radiation (with opening angle $\theta_j = 0.1$) is a very powerful gamma energy radiation with typical emitted energy in the range of 10^{51} erg lasting a few tens of seconds and with a peak photon energy E_p of a few hundreds of keV. The compact nature of the GRB progenitors and the observed photon energy spectra most likely require both photon propagation transparency and relativistic motion of the emitting source. The evolution of a typical GRB can be for simplicity divided in two different phases: the first phase occurs close to the progenitor and energizes and collimates the emitting particles. This first phase may be related to the emission of quasi thermal or Comptonized photons (Meszaros & Rees, 2005). A second step can lead to efficient particle acceleration far out from the central source, and eventually to efficient radiation by synchrotron emission in internal and/or external shocks.

In this paper, we investigate a model of GRB production and evolution that assumes: (1) the existence of a thermal-like precursor of very large intensity following the energy release from relativistically moving shells or hydrodynamic fronts originating from the GRB compact object; (2) the radiative interaction of such a precursor (with a given power-spectrum as seen by an observer at rest) with a surrounding (initially not-ionized) gas shell at rest of density $n \sim 10^9 \mathrm{cm}^{-3}$ at the approximate distance $R \sim 10^{15}$ cm; (3) propagation of the radiative front within the shell, with possible activation of wake-plasma particle acceleration at the outwardly propagating radiative shock; (4) wake-field particle acceleration producing a generation of non-thermal electrons whose kinetic energy distribution function extends up to $> 100\,\mathrm{MeV}$; (5) betatron radiation of these energized electrons producing the observed non-thermal prompt GRB high-energy emission in the X-ray/gamma-ray energy ranges. The existence of an amount of material corresponding to a Thompson optical depth 1 is derived by the observation made by BATSE of late bumps in the GRB light curve (Connaughton 2002), interpreted as a Compton reprocessed emission from dense circum-burst material at $R = 10^{15}$ cm (Barbiellini et al., 2004). The physical origin of this material is at present subject of very intense simulations of the WR wind, where the wind could be compressed by the preceding supernova explosion. In this paper we would like to consider only the experimental evidence. In this scenario, we assume that the energy attributed to the precursor (or to any electromagnetic signal entering the plasma) is E_γ/ϵ where E_γ is the energy in the prompt emission and ϵ is the conversion efficiency of the precursor energy in the prompt emitted radiation. Since the photon component of the precursor has quasi-thermal energy, the absorption is very large; the highest energy component of the prompt is absorbed at a level of 70%. Our GRB emission model does not imply a relevant role for the intrinsic or MHD-driven comoving magnetic field in the ejecta and/or radiative fronts. From this point of view, it provides an emission scenario quite different from those assumed in current theoretical models of prompt GRB emission such as the synchrotron shock model (Tavani 1996; Lloyd & Petrosian 2002), the jitter radiation model (Medvedev 2000), the diamagnetic relativistic pulse accelerator (Liang et al. 2003), and the saturated Comptonization model (e.g., Liang et al. 1997).

The very recent technical progress on high power laser beams has provided experimental evidence supporting the concept of wake-field particle acceleration in a plasma excited by an electromagnetic wave of duration comparable or shorter than the inverse of the plasma frequency (Tajima and Dawson 1979). A laser beam with a duration of 30 femtoseconds and a power of 50 TeraWatts, illuminating a gas jet 1 mm long with a density of 10^{19} particles per cm^3 is able to accelerate electrons to an energy of few tens of MeV. An X-ray beam of a few keV generated by the electron radiation loss in the plasma was observed by Ta Phuoc et al. (2005) with a characteristic angular distribution resembling that of synchrotron radiation (this motivates the qualification used in the literature of “Laser driven synchrotron radiation”). Previous connections between the laboratory plasma physics and the astrophysical plasmas have been suggested by Chen et al. (2002), which proposed a wake field acceleration excited by the Alfven shocks, producing Ultra High Energy Cosmic Rays (UHECR). Simulations using three dimensional particle-in-cell code, have been recently done by Ng & Noble (2005) studying the dynamics of a neutral, narrow, electron-positron jet propagating through an unmagnetized electron-ion plasma. In this paper, following the general hypothesis already suggested by Tajima & Dawson (1979), the processes of acceleration and radiative losses induced by ultra short laser pulses on the gas target, validated by recent experimental data, are suggested to be the source of the GRB prompt emission. In our case, the laser role is played by GRB precursor photons emitted by the γ and $e^\pm$ shells collimated in an angle θ_j and moving with high relativistic velocity and the plasma density of the target gaseous nebula is assumed to be on the average 10^9 cm^{-3} over the region producing the prompt emission at a distance from the central source of about 10^{15} cm.

In order to activate the coherent wake-field particle acceleration in the target plasma in the laboratory, where $n_{e0} \sim 10^{19}$ cm^{-3}, the threshold power surface density is

$$\sigma_{Wt} \sim 10^{18} \mathrm{W cm}^{-2}$$

Furthermore, assuming a coherent precursor emission the two conditions need to be satisfied: (1) the radiation has to be suddenly delivered (in the form of pulses or micro-bursts) to the target gas within ultra-short timescales τ_p (that for our conditions need to satisfy the condition $\tau_p < 10^{-9}$ s in the observer/nebula frame); and (2) the radiation intensity delivered within these ultra-short pulses has to be sufficiently large to activate microscopic plasma electron-ion separation. In the following, we assume that the GRB initial fireball producing a very intense radiative precursor is satisfying condition (2). In particular, this is the key point in the analogy GRB-Lab plasma acceleration is that the coherent radiation of laboratory lasers can be translated into non-coherent (stochastic) radiation from the precursor if a price is payed in terms of power density. The critical parameter in the coherent Lab plasma acceleration is the power surface density threshold. We show below that σ_{Wt} scales linearly with particle density. Its value is assumed from the experiment:

$$\sigma_{Wt} = 3 \times 10^{18} \left(\frac{n}{10^{19}\ cm^{-3}} \right) \mathrm{W\ cm}^{-2} \tag{1}$$

In the hypothesis that the GRB precursor carries an energy $E_{\mathrm{pr}} = E_\gamma/\epsilon$ and assuming $t_{\mathrm{pr}} \sim t_{grb} \sim 30/(1+z)$ s the power density on the surface $R^2\theta^2 = 10^{27}$ cm^2 is:

$$\sigma_{Wgrb} = \frac{E_{\mathrm{pr}}}{t_{grb} R^2 \theta^2} \geqslant 2 \times 10^{16}\ \mathrm{W\ cm}^{-2} \tag{2}$$

which exceeds by several order of magnitude (around 10^8) the power density threshold (scaled at $n = 10^9$ cm^{-3}).

The threshold to initiate stochastically the wakefield mechanism is equal to the coherent power density multiplied by the factor $\sqrt{t_{\mathrm{pr}}/\tau_p}$ that is is the probability to produce by chance the coherent WFA, independently in time and space.

In other words, GRB not-coherent prompt emission is much more powerful than the coherent lab radiation: we assume that the loss of coherence is balanced by the enormous amount of power per square centimeter as well as the possibility of stochastic resonance: the wake field acceleration threshold is overcome.

Figure 1 from Ta Phuoc et al. (2005) shows schematically the photon emission principle. If the plasma is excited by the precursor photons, the excited region and the emission of the radiated photons are likely to be in a cone of angular aperture equal to θ_j starting at a distance of 10^{15} cm from the central compact object.

Stochastic acceleration and deceleration are in principle related by plasma effects. However, since they are statistically independent, the two processes can be separated without loss of generality. In the following, the acceleration phase is not treated in detail. On the basis of the experimental results of Ta Phuoc et al. (2005) it is assumed that a power law energy spectrum of electron and positron is produced. The accelerated beam enters in the plasma region with constant gas density and radiates in the plasma, following again the results of Ta Phuoc et al. (2005).

The value of the plasma density in the GRB environment is assumed following Barbiellini et al. (2004). Because of the relation among the precursor and the radiating electrons, to reproduce the GRB data a stochastic treatment of the deterministic formulae of Ta Phuoc et al. (2005) is applied.

In the laser-driven betatron radiation the emission follows the scheme of Ta Phuoc et al. (2005)[fig. 1]. The angular distribution of the emitted radiation is dominated by the wiggling angle of the electron,

$$\theta = \frac{K}{\gamma}$$

where

$$K = \frac{\gamma \omega_b}{c} r_0$$

with

$$\omega_b = \frac{\omega_p}{\sqrt{2\gamma}}$$

the oscillating frequency, where $\omega_p = \sqrt{ne^2/(\epsilon_0 m_e)}$ is the plasma fundamental frequency and r_0 that could be estimated as:

$$r_0 = \frac{c}{\omega_p \sqrt{\gamma}} = \frac{\bar{\lambda}_p}{\sqrt{\gamma}}$$

is equivalent to the impact parameter of the electron into the positive plasma column (e is the electron's charge). Interpreting $\bar{\lambda}_p = c/\omega_p$ as the screen distance of a charge inside the plasma, if the charge is moving relativistically $\bar{\lambda}_b = c/\omega_b$ and r_0 represent respectively the longitudinal and transverse screening distance.

2 SCALING RELATIONS IN COHERENT PLASMA ACCELERATION

The key parameter in the coherent wakefield acceleration (CWFA) is the power surface density σ_W needed to produce the complete charge separation over a lenght $\bar{\lambda}_p$ on a timescale $\tau_p = \omega_p^{-1}$.

The energy volume density ρ_ϵ for complete separation could be espressed as:

$$\rho_\epsilon = E^2 \epsilon_0 = \frac{\sigma^2}{\epsilon_0} = \frac{e^2 n^2 \bar{\lambda}_p^2}{\epsilon_0}$$

where σ is the charge surface density. The quantity ρ_ϵ scales linearly with n while the energy surface density $\sigma_\epsilon = \rho_\epsilon \bar{\lambda}_p$ varies as:

$$\sigma_\epsilon = \frac{e^2 n^2 \bar{\lambda}_p^3}{\epsilon_0} \propto n^{1/2}.$$

The key parameter $\sigma_W = \sigma_\epsilon \omega_p$ then scales as the plasma particle density n.

Using the relation among θ and the plasma quantities $\theta = \frac{r_0}{\bar{\lambda}_b} \sim 1/\gamma$, since $r_0 = \bar{\lambda}_p / \sqrt{\gamma}$ it is possible to derive:

$$\gamma r_0^2 \propto n^{-1} \tag{3}$$

Let us consider an electron radiating with the typical formula:

$$\frac{dE}{dt} = -m\, c^2 \frac{d\gamma}{dt} = \frac{e^2 \gamma^4}{6\pi\epsilon_0 c^3} |a_\perp|^2 \tag{4}$$

where $a_\perp$ is the transverse electron acceleration

$$a_\perp = r_0 \omega_b^2 = r_0 \frac{\omega_p^2}{2\gamma}.$$

The energy loss by the electron in one oscillating period is:

$$\frac{d\gamma}{dt} \propto \gamma^4 r_0^2 \omega_b^4$$

with $\Delta t = \omega_b^{-1}$. At the equilibrium $\Delta\gamma \propto \gamma^{2.5} r_0^2 n^{3/2} = \text{const}$. Recalling Eq.3, we obtain:

$$\gamma^{3/2} n^{1/2} = \text{const}$$

and, hence, $\gamma \propto n^{-1/3}$. Furthermore, using again the relation $\gamma r_0^2 \propto n^{-1}$ it is also possible to derive that r_0 scales as $n^{-1/3}$. Another way to obtain the previous relation is to consider that the restoring factor K does not depend on the density n so that:

$$K \propto \omega_p\, \gamma^{1/2}\, r_0 \;\propto n^{1/2}\, n^{-1/6}\, r_0 = \text{const}$$

obtaining that $r_0 \propto n^{-1/3}$.

Notice that the maximum electron Lorentz factor $\gamma_{\max}$ measured by Ta Phuoc et al. (2005) is, roughly $\gamma_{\max} \sim 200$. For a density of 10^9 cm^{-3}, the electrons can be accelerated up to:

$$\gamma_{\max} \sim 200 \left(\frac{10^{19}}{10^9}\right)^{1/3} \sim 4.3 \times 10^5$$

In case of low densities, wake field acceleration can be as effective as producing high energy cosmic rays and, consequently gamma rays.

Following Ta Phuoc et al. (2005) and using the fundamental frequency of radiation ω_f in the case the plasma acts as a wiggler we have

$$\omega_f = \frac{3}{2} \gamma^3 c r_0 \frac{\omega_p^2}{2\gamma c^2} = \frac{3}{4} c \gamma^2 \frac{r_0}{\bar{\lambda}_p^2}$$

.

We obtain an estimate of the peak energy of the emitted radiation E_{peak}:

$$E_{\text{peak}} = \hbar\omega_f = \frac{3}{4} \hbar c \gamma^2 \frac{r_0}{\bar{\lambda}_p^2}$$

The energy distribution of the accelerated electrons is

$$N(\gamma) = N_0 \gamma^{-2}$$

assuming the power law index again from the experiment (Ta Phuoc et al. 2005).

The peak energy computed as an average on the electron spectrum is then:

$$\langle E_{\text{peak}} \rangle = \frac{3}{4} \hbar c \frac{r_0}{\bar{\lambda}_p^2} \gamma_{\min} \gamma_{\max}$$

3 GRB GAMMA EMISSION IN THE STOCHASTIC WAKE FIELD ACCELERATION REGIME

We address here the GRB prompt radiation produced by electrons radiating in a plasma excited by a very intense precursor playing the role of a "laser". The basic formulae of laser-driven betatron radiation are assumed taking into account the stochastic nature of the acceleration and the radiation phases. Let us now apply these results to the case of a plasma accelerated by the GRB precursor photons. We will treat the motion of the relativistic electrons radiating in the plasma. The fundamental difference with the above treatment resides in the stochastic nature of acceleration and radiation in the case of the GRB emission.

If the power surface density σ_W at any position of the electrons producing the GRB exceeds the value measured by the experiment (Ta Phuoc et al. 2005) scaled at the GRB density and multiplied by the stochastic factor, the radiated photons of the GRB will be emitted from any accelerated electron provided that the stochastic angle of the electrons θ_{eff} remain below the jet opening angle θ_j. The elementary angle of the electron oscillation is:

$$\theta_i = \frac{r_0}{\bar{\lambda}_b}$$

The stochastic motion of the electrons in the plasma results in an effective solid angle spanned by the electrons:

$$\theta_{\text{eff}}^2 = \left(\frac{R}{\bar{\lambda}_b}\right) \theta_i^2$$

where we have introduced the adimensional quantity N:

$$N = \left(\sqrt{\frac{R}{\bar{\lambda}_b}}\right) = \sqrt{\frac{R}{\sqrt{2\,\gamma}\,\bar{\lambda}_p}}$$

where R is the distance travelled by the electron inside the excited plasma. This will be used to estimate the stochasticity of the electron geometrical path.

An electron with energy $E = \gamma m c^2$ wiggling inside the plasma with density n will contribute to the GRB detectable emission for a path R_θ if:

$$[\theta_{\rm eff}(R_\theta, \gamma, n)]^2 \leqslant \theta_j^2 \tag{5}$$

Putting esplicitly the dependence on the path R_θ the previous equation could be written as:

$$R_\theta \frac{r_0^2}{\lambda_p^3}(2\gamma)^{-1.5} \leqslant \theta_j^2$$

This relation expressed in terms of the plasma density and using the scaling laws for r_0, λ_p and γ derived previously could be written as:

$$R_\theta(\gamma) = 2\sqrt{2}\lambda_p \theta_j^2 \gamma^{5/2}$$

The equation 5 introduces a threshold for the electron emission over a distance R. The threshold Lorentz factor γ_t is obtained requiring that even in the case of a large energy loss, the electron trajectory still remains within the energized cone.

The effective energy loss in the stochastic acceleration regime is:

$$\frac{d\gamma}{dt} = -\frac{2}{3} r_e r_0^2 \gamma^4 \frac{\omega_b^4}{c^3}\left(\frac{R}{\lambda}\right)^{0.5}$$

where for simplicity we have introduced the classical electron radius r_e

$$r_e = \frac{e^2}{4\pi\epsilon_0 m_e c^2}$$

and where we have multiplied the coherent energy loss by the stochastic factor $\sqrt{R/\lambda}$. This factor is our best estimation of the effect of the turbulence of the circumburst medium. Expressing the previous relation in terms of γ, and substituting $dt = c\, dR$, we obtain:

$$d\gamma\gamma^{-3/4} = \frac{1}{11} r_e \lambda_p^{-5/2} dR^{3/2}$$

$$\gamma^{1/4} = \frac{r_e}{8.3}\lambda_p^{-5/2} R^{3/2}$$

From this relation we estimate a typical "energy loss" distance R_e equivalent to the path where the electron looses half of its energy. Integrating the previous equation we get:

$$R_e \sim 4.1\, \lambda_p \left(\frac{\lambda_p}{r_e}\right)^{2/3} \gamma^{1/6}$$

and

$$\frac{R_e}{\lambda_b} = 2.8\, \gamma^{-1/3}\left(\frac{\lambda_p}{r_e}\right)^{2/3}$$

The value of γ_t is then obtained by requiring

$$R_\theta(\gamma_t) = R_e(\gamma_t)$$

obtaining

$$\gamma_t = 1.2\left(\frac{\lambda_p}{r_e}\right)^{2/7} \theta_j^{-6/7}$$

The opening angle of the jet θ_j imposes a threshold on the energy of the emitting electron wiggling with an effective angle so that the mean value of the peak energy has to be computed over the electron spectrum integrated only between γ_t and $\gamma_{\rm max}$.

Taking into account the stochastic nature of the phenomenum due to the external density turbolence, we can calculate the predicted value for $\langle E_{\rm peak}\rangle_{\rm eff}$ in the stochastic regime, simply multiplying the coherent value of $\langle E_{\rm peak}\rangle$ by the stochastic factor $\sqrt{R_e/\lambda_b}$. After an integration between $2\gamma_t$ and γ_t, assuming that only these electrons contribute to the observed radiation, we obtain:

$$\langle E_{\rm peak}\rangle_{\rm eff} = 2.5\frac{\hbar c}{\lambda_p}\left(\frac{\lambda_p}{r_e}\right)^{1/3}\gamma_t^{4/3}$$

$$\langle E_{\rm peak}\rangle_{\rm eff} = 3.0\frac{\hbar c}{\lambda_p}\left(\frac{\lambda_p}{r_e}\right)^{5/7}\theta_j^{-8/7}$$

From the previous relations, we have:

$$\langle E_{\rm peak}\rangle_{\rm eff} \approx 25\ {\rm keV}\frac{\rm n}{10^9}\theta_{\rm j}^{-8/7}$$

Assuming the density $n = 10^9\ {\rm cm}^{-3}$ obtained by the analysis of the Compton tail phenomenum (Barbiellini et al. 2004), and θ_j=0.1 ($\theta_j \approx 6°$) we obtain in the GRB source frame:

$$\langle E_{\rm peak}\rangle_{\rm eff} \approx 350\ {\rm keV}$$

4 SPECTRAL - ENERGY CORRELATIONS

Noticing that the quantity γ_t introduces then a dependence of $\langle E_{\rm peak}\rangle$ on θ_j, we use the above derived formulae for the peak energy of the emission and the effective angle of the radiation to derive an estimate for the recently obtained Amati (Amati et al., 2002) and Ghirlanda (Nava et al. 2006) relations.

Considering the electrons with an energy spectrum similar to that derived experimentally in the coherence regime, with Lorentz factor between γ_t and $2\gamma_t$, the mean useful electron energy is:

$$E_\gamma = N_0 mc^2 \frac{\int \gamma\gamma^{-2} d\gamma}{\int \gamma^{-2} d\gamma} = 2\ln 2 N_0 mc^2 \gamma_t$$

If each electron of energy greater than $mc^2\gamma_t$ emit n_γ of mean energy $\langle E_{\rm peak}\rangle$ then the energy conservations requires:

$$mc^2\langle\gamma\rangle = n_\gamma \langle E_{\rm peak}\rangle$$

$$(2\ln 2)\, mc^2\gamma_t = n_\gamma \frac{\hbar c}{\lambda_p}\left(\frac{\lambda_p}{r_e}\right)^{1/3}\gamma_t^{4/3}$$

The energy conservation requires $n_\gamma \propto \gamma_t^{-1/3} \propto \langle E_{\rm peak}\rangle^{-1/4}$. The total energy E_γ being composed of the emission by the N_0 electrons results

$$N_0 mc^2\langle\gamma\rangle = E_\gamma = N_0 n_\gamma \langle E_{\rm peak}\rangle = N_0 n_\gamma \langle E_{\rm peak}\rangle^{1/4}\langle E_{\rm peak}\rangle^{3/4}$$

Obtaining:

$$E_\gamma = N_0 \times {\rm const} \times \langle {\rm E_{peak}}\rangle^{3/4}$$

If the isotropic energy $E_{\rm iso}$ is introduced:

$$E_{\rm iso} \approx \frac{E_\gamma}{\theta_j^2} = \frac{N_0 n_\gamma \langle E_{\rm peak}\rangle}{\theta_j^2} \propto \theta_j^{-2}\langle E_{\rm peak}\rangle^{3/4} \propto \langle E_{\rm peak}\rangle^{5/2}$$

Inverting the previous equation we obtain $E_{\rm peak} \propto E_{\rm iso}^{0.4}$ as outlined in the so called "Amati" relation.

5 CONCLUSION

We studied the properties of a novel mechanism for prompt GRB radiation based on plasma laser-wake acceleration and radiation induced by an intense precursor irradiating a static gas nebula. The stochastic radiative emission of the energized plasma produces a photon emission with angular distribution dominated by the electron trajectories. Under general conditions applicable to our GRB model (precursor, static cloud, laser-wake acceleration conditions) we showed that there is a relation between the observed jet property (θ_j) and an electron energy threshold. In our mechanism the energy of the emitting wiggling electrons is the typical energy of the laser-wake betatron radiation. Furthermore, we derived a relation between source properties such as the collimation angle and the emitted energy.

ACKNOWLEDGEMENTS

We would like to thank Ugo Amaldi for the suggestion to investigate the plasma wake field acceleration as an efficient mechansim, Victor Malka and Danilo Giuletti for the illustration of the fundamental elements on plasma acceleration. One of us (G.B.) would like also to express gratitude to Giuseppe Cocconi for the wise suggestion on how to deal with the comparison of events taking place within environments with scale different by many order of magnitude.

REFERENCES

L. Amati et al., *A&A*, **390**, 81 (2002).
G. Barbiellini et al., *MNRAS*, **350**, L5 (2004).
P. Chen, T. Tajima, and Y. Takahashi, *Phys. Rev. Lett.*, **89**, 161101 (2002).
V. Connaughton, *ApJ*, **567**, 102 (2002).
E. Liang, M. Kukunose, I.A. Smith, and A. Crider, *ApJL*,**479**, L35 (1997).
E. Liang, K. Nishimura, H. Li, and S.P. Gary, *Phys. Rev. Lett.*, **90**, 085001 (2003).
N.M. Lloyd-Ronning, and V. Petrosian, *ApJ*, **565**, 182 (2002).
M.V. Medvedev, *ApJ*, **540**, 704 (2000).
L. Nava et al., `astro-ph 0511499`, *A&A* accepted (2006)
J. S. T. Ng, & R. J. Noble, preprint astro-ph/0507231.
T.Tajima and J.M. Dawson, *Phys. Rev. Lett.*, **43**, 267 (1979).
K. Ta Phuoc, et al., *Physics of Plasmas*, **12**, 023101 (2005).
M. Tavani, *Phys. Rev. Lett.*, **76**, 3478 (1996).
M.J. Rees and P. Meszaros, *ApJ*, **628**, 847 (2005)